

COMPUTING THE EXPONENTIAL MAP ON $\mathrm{SO}(3)$

ADAM WILLIAMS

1. OVERVIEW

This document details how to compute various quantities related to the Lie group $\mathrm{SO}(3)$ and its Lie algebra $\mathfrak{so}(3)$, for practical engineering purposes. Recall that $\mathrm{SO}(3)$ is the set of orthogonal, orientation-preserving linear transformations on \mathbb{R}^3 and can be realized as a matrix group:

$$\mathrm{SO}(3) = \{R \in \mathrm{GL}(3) \mid RR^\top = I\}.$$

2. CHARACTERIZING THE LIE ALGEBRA $\mathfrak{so}(3)$

2.1. Skew-Symmetric Matrix Representation. Differentiating a curve R in $\mathrm{SO}(3)$ with $R(0) = I$ yields

$$R'(0)^\top + R'(0) = 0,$$

meaning that the Lie algebra $\mathfrak{so}(3)$ can be realized as the vector space of 3×3 skew-symmetric matrices, with Lie bracket given (as with all matrix groups) by the commutator: $[X, Y] = XY - YX$. Hence we can choose the basis $\mathcal{G} = (G_1, G_2, G_3)$

$$\begin{aligned} G_1 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \\ G_2 &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \\ G_3 &= \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \end{aligned}$$

as a choice of exponential coordinates (or infinitesimal generators) on $\mathrm{SO}(3)$, and do computations with respect to this basis.

2.2. Axis-Angle Vector Representation. An alternative realization of $\mathfrak{so}(3)$ is the vector space \mathbb{R}^3 with Lie bracket given by the cross product:

$$[\omega, \eta] = \omega \times \eta.$$

The isomorphism is given by the mapping

$$\begin{aligned}\mathbb{R}^3 &\cong \mathfrak{so}(3) \\ \omega &\mapsto \omega_{\times}\end{aligned}$$

taking $\omega \in \mathbb{R}^3$ to its *skew matrix*

$$\omega_{\times} = \begin{pmatrix} 0 & -\omega_z & \omega_y \\ \omega_z & 0 & -\omega_x \\ -\omega_y & \omega_x & 0 \end{pmatrix} = \omega_x G_1 + \omega_y G_2 + \omega_z G_3.$$

The **axis-angle** vector ω has the following geometric interpretation: Let

$$\begin{aligned}\hat{\omega} &= \omega / \|\omega\| \\ \theta &= \|\omega\|.\end{aligned}$$

Then $\exp(\omega)$ is a right-handed rotation of angle θ around the axis $\hat{\omega} \in \mathbb{R}^3$. Hence the infinitesimal generators G_1 , G_2 , and G_3 that we have chosen are perturbations around the x , y , and z axis, respectively. And the one-parameter subgroups $\theta \mapsto \exp(\theta\hat{\omega})$ compound infinitesimal rotations around a given axis for an angle of θ .

3. EXPONENTIAL AND LOGARITHM

3.1. Exponential. The goal of this section is to compute the exponential map, taking an axis-angle vector to its rotation matrix in $\text{SO}(3)$, using the matrix exponential:

$$\exp(\omega) = \sum_{k=0}^{\infty} \frac{\omega_{\times}^k}{k!}.$$

Let $\hat{\omega} = \omega / \|\omega\|$, and $\theta = \|\omega\|$. Using properties of the cross product, we have $\hat{\omega}_{\times}^3 = -\hat{\omega}_{\times}$, and so higher powers of $\hat{\omega}_{\times}$ in the power series collapse:

$$\begin{aligned}\hat{\omega}_{\times}^3 &= -\hat{\omega}_{\times} \\ \hat{\omega}_{\times}^4 &= -\hat{\omega}_{\times}^2 \\ \hat{\omega}_{\times}^5 &= \hat{\omega}_{\times}.\end{aligned}$$

Hence we can group all of the power series terms into a $\hat{\omega}_{\times}$ term and a $\hat{\omega}_{\times}^2$ term:

$$\begin{aligned}\exp(\omega) &= I + \left(1 - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} + \cdots\right) \hat{\omega}_{\times} + \left(\frac{\theta^2}{2!} - \frac{\theta^4}{4!} + \frac{\theta^6}{6!}\right) \hat{\omega}_{\times}^2 \\ &= I + \left(\frac{\sin \theta}{\theta}\right) \omega_{\times} + \left(\frac{1 - \cos \theta}{\theta^2}\right) \omega_{\times}^2.\end{aligned}$$

Hence we have a practical formula for the exponential of a skew matrix. Note that, in practice, the trigonometric coefficients should be computed using a Taylor series

approximation for sufficiently small θ :

$$\begin{aligned}\frac{\sin(\theta)}{\theta} &= 1 - \frac{\theta^2}{6} + \frac{\theta^4}{120} + O(\theta^6) \\ \frac{1 - \cos \theta}{\theta^2} &= \frac{1}{2} - \frac{\theta^2}{24} + \frac{\theta^4}{720} + O(\theta^6)\end{aligned}$$

3.2. Logarithm. In order to derive a formula for the (partial) inverse of \exp , we must

- (1) Determine the **normal neighborhood** $U \ni 0 \in \mathfrak{so}(3)$ on which \exp is invertible.
- (2) Compute a formula for $\log : \exp(U) \cong U$.

First, to compute the angle of rotation $\theta = \|\omega\|$. If we take the trace of $R = \exp(\omega)$, then applying linearity and the fact that ω_{\times} is traceless, we obtain

$$\begin{aligned}\text{tr } R &= \text{tr } I + 0 + \left(\frac{1 - \cos \theta}{\theta^2} \right) \text{tr } \omega_{\times}^2 \\ &= 3 + \left(\frac{1 - \cos \theta}{\theta^2} \right) (-2\|\omega\|)\end{aligned}$$

and thus

$$\cos \theta = \frac{\text{tr } R - 1}{2}.$$

Restricting $\theta \in [0, \pi]$ then gives the unique solution

$$\theta = \arccos \frac{\text{tr } R - 1}{2}.$$

Now to compute the axis of rotation. we first recall the fact that the set of $n \times n$ matrices $M_n(\mathbb{R})$ is the direct (vector space) sum of the set of skew-symmetric matrices and the set of symmetric matrices:

$$M_n(\mathbb{R}) = \text{Skew}_n(\mathbb{R}) \oplus \text{Sym}_n(\mathbb{R}).$$

That is, any square matrix can be written *uniquely* as a skew-symmetric matrix and a symmetric matrix. In fact, this decomposition has a simple formula:

$$A = \frac{A - A^{\top}}{2} + \frac{A + A^{\top}}{2}.$$

Now, let $R = \exp(\omega)$. Observe that the formula for \exp from the previous section is actually the skew-symmetric/symmetric decomposition of R :

$$R = \underbrace{\left(\left(\frac{\sin \theta}{\theta} \right) \omega_{\times} \right)}_{\frac{R - R^{\top}}{2}} + \underbrace{\left(I + \left(\frac{1 - \cos \theta}{\theta^2} \right) \omega_{\times}^2 \right)}_{\frac{R + R^{\top}}{2}}.$$

Hence, if $\sin \theta \neq 0$, we can use the skew-symmetric part to solve for ω_\times :

$$\omega_\times = \left(\frac{\theta}{2 \sin \theta} \right) (R - R^\top).$$

And so we simply choose ω such that

$$\omega_\times = R - R^\top.$$

Note that, when θ is small, it is best to compute the coefficient using a Taylor series:

$$\frac{\theta}{2 \sin \theta} = \frac{1}{2} + \frac{\theta^2}{12} + \frac{7\theta^4}{720} + O(\theta^6).$$

Now, this method breaks down when $\theta > 0$ and $\sin(\theta) = \pi$ (note that if $\theta = 0$ then we choose $\omega = 0$, trivially). So, we have a unique log for all ω within the interior of a ball of radius π . At the boundary $\theta = \pi$, however, R is purely symmetric (hence $R = R^\top = R^{-1}$) and so the skew matrix provides no information. Note that this is geometrically intuitive: A rotation of angle π is its own inverse. Hence we must use the symmetric part to solve for $\pm\omega$, and make an arbitrary choice of sign: setting $\theta = \pi$, we have

$$R = I + 2\hat{\omega}_\times^2.$$

Note that $\hat{\omega}_\times^2 = \hat{\omega}\hat{\omega}^\top - I$, and hence

$$\hat{\omega}\hat{\omega}^\top = \frac{1}{2}(R + I).$$

we thus have

$$\begin{aligned} \hat{\omega}_x^2 &= \frac{1}{2}(r_{11} + 1) \\ \hat{\omega}_y^2 &= \frac{1}{2}(r_{22} + 1) \\ \hat{\omega}_z^2 &= \frac{1}{2}(r_{33} + 1), \end{aligned}$$

and the relative signs of the entries of $\hat{\omega}$ can be determined by the off-diagonal components. This determines ω up to a factor of ± 1 .

3.3. Adjoint Representation of the Group. The adjoint representation of a group element $R \in \text{SO}(3)$ is the linear mapping

$$\omega_\times \mapsto R\omega_\times R^\top.$$

from $\mathfrak{so}(3) \rightarrow \mathfrak{so}(3)$. We would like to write this linear transformation with respect to the basis \mathcal{G} , i.e. express it as a left multiplication by a 3×3 matrix

$$\omega \mapsto \text{Ad}_R \omega.$$

This is straightforward, using the fact that the cross product is preserved by rotation: $Rx \times Ry = R(x \times y)$. Let $y \in \mathbb{R}^3$ be arbitrary, and observe

$$\begin{aligned} [\text{Ad}_R \omega]_{\times} Ry &= [R\omega_{\times} R^{\top}] Ry \\ &= R(\omega \times y) \\ &= R\omega \times Ry \\ &= [R\omega]_{\times} Ry. \end{aligned}$$

Hence $[\text{Ad}_R \omega]_{\times} = [R\omega]_{\times}$, meaning

$$\text{Ad}_R = R.$$

We say that $\text{SO}(3)$ is *self-adjoint*.

3.4. Adjoint Representation of the Algebra. The adjoint representation of a Lie algebra element $\omega \in \mathfrak{so}(3)$ is the linear mapping

$$\eta_{\times} \mapsto \omega_{\times} \eta_{\times} - \eta_{\times} \omega_{\times}$$

from $\mathfrak{so}(3) \rightarrow \mathfrak{so}(3)$. We can derive the matrix for this mapping with respect to \mathcal{G} by applying the *Jacobi identity* for the cross product: For any $x \in \mathbb{R}^3$, and any vectors $\omega, \eta \in \mathfrak{so}(3)$,

$$\omega \times (\eta \times x) + \eta \times (x \times \omega) + x \times (\omega \times \eta) = 0.$$

Applying anti-commutativity,

$$\omega \times (\eta \times x) - \eta \times (\omega \times x) = (\omega \times \eta) \times x.$$

Hence

$$\begin{aligned} [\text{ad}_{\omega} \eta]_{\times} x &= \omega_{\times} \eta_{\times} x - \eta_{\times} \omega_{\times} x \\ &= \omega \times (\eta \times x) - \eta \times (\omega \times x) \\ &= (\omega \times \eta) \times x \\ &= [\omega \times \eta]_{\times} x. \end{aligned}$$

Hence $[\text{ad}_{\omega} \eta]_{\times} = [\omega \times \eta]_{\times} = [\omega_{\times} \eta]_{\times}$, meaning

$$\text{ad}_{\omega} = \omega_{\times}.$$